

Environmental Information Sheet

BROADWAY STAR MAFF/MAPP 14319



A water dispersible granule formulation containing 7.08%w/w pyroxsulam and 1.42%w/w florasulam, triazolopyrimidine herbicides (plus 7.08%w/w cloquintocet-mexyl as a safener) for the control of weeds in winter wheat.

Maximum application rate: 265 g product/ha

Maximum one application per crop

Section	Profile
<p>1. WILDLIFE</p> <p>Mammals and Birds</p>	<p>Broadway Star is not classified as <i>"Harmful to game and wildlife"</i></p> <p>No risk management is necessary to protect mammals and birds.</p> <p>Broadway Star has a low toxicity to mammals and birds. It poses a low risk to birds and mammals living and feeding in the treated area.</p>
<p>2. BEES</p>	<p>No risk management is necessary. There is no requirement to avoid application of Broadway Star when bees may be foraging on flowering weeds.</p>
<p>3. NON TARGET INSECTS AND OTHER ARTHROPODS</p>	<p>No risk management is necessary. Broadway Star poses a low risk to arthropod species found in and around treated fields, such as predatory mites and parasitic wasps.</p>
<p>4. AQUATIC LIFE</p>	<p>Broadway Star is classified <i>"VERY TOXIC TO AQUATIC ORGANISMS, may cause long-term adverse effects in the aquatic environment"</i></p> <p>Broadway Star has a moderate toxicity to fish, a low toxicity to <i>Daphnia</i> (water fleas) but is extremely toxic to algae and the aquatic plant <i>Lemna</i>. Care must be taken to ensure that surface waters or ditches are not contaminated with the product or the used containers.</p> <p>Risk management is essential. To protect aquatic organisms <i>"DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1m of the top of a ditch which is dry at the time of application. Aim spray away from water"</i>.</p>

Environmental Information Sheet

BROADWAY STAR MAFF/MAPP 14319



Section	Profile
<p>5. SOIL and GROUNDWATER</p> <p>Earthworms</p> <p>Soil Micro-organisms</p>	<p>Pyroxsulam and florasulam have a low persistence in soil.</p> <p>Under laboratory conditions they are weakly adsorbed and considered to be highly mobile. However, studies under field conditions and detailed modelling have demonstrated that the low persistence, coupled with very low application rates, prevents significant groundwater contamination.</p> <p>Broadway Star has a low toxicity to earthworms. The risk to earthworms is low and no risk management is necessary.</p> <p>No risk Management is necessary. At up to at least 5 times the recommended application rate Broadway Star has negligible effects on soil microbial respiration and nitrogen turnover. The risk to soil microbial activity is therefore low.</p>
<p>6. NON-TARGET PLANTS</p>	<p>Broadway Star is a post emergence herbicide.</p> <p><i>“Extreme care must be taken to avoid spray drift onto susceptible crops and non-crop plants outside the target area”.</i></p>

ALWAYS READ THE LABEL: USE PESTICIDES SAFELY

Care must be taken to minimise the risk of surface water contamination from farmyard and field sources.

For further information about the environmental profile of this product contact:-

Company: Dow AgroSciences Limited
 Address: Latchmore Court, Brand Street, Hitchin, Hertfordshire. SG5 1NH
 Telephone: 01462 457272
 Fax: 01462 426605
 e-mail: ukhotline@dow.com

24 Hour Emergency Telephone Number +44 (0) 1553 761 251

This Environmental Information Sheet was prepared in accordance with CPA Guidance Notes Version 4.

© copyright of Dow AgroSciences Limited