BlueN™

NUTRIENT EFFICIENCY BIOSTIMULANT

BIOSTIMULANT TO INCREASE NUTRIENT EFFICIENCY, UPTAKE AND AVAILABILITY.

Composition:

Methylobacterium symbioticum SB23, 3 x 10⁷ CFU/g Wettable Powder

TO AVOID RISKS TO HUMAN HEALTH AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE

Registration holder: Symborg Business Development S.L.

Avda. Jesús Martínez Cortado, 51. P.I. Cabezo Cortado, 30100, Murcia,

Spain

Manufactured by: Symborg Business Development S.L.

Avda. Jesús Martínez Cortado, 51. P.I. Cabezo Cortado, 30100, Murcia,

Spain

Distributed by: Corteva Agriscience UK Limited, CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE,

Telephone: +44 (0)1462) 457272

24 Hour Emergency Telephone Number: 076 680 5288

National Poison Centre (Dublin): 01 809 2166

Technical Hotline: +44 (0)800 689 8899, e-mail: ukhotline@corteva.com

Lot Number: See packaging Manufacturing date: See packaging Expiration Date: See packaging



Powered by Symborg

Precautionary statements

Follow the instructions for use to avoid risks to human health and the environment.

- P102: Keep out of reach of children.
- P261: Avoid breathing dust.
- P262: Do not get in eyes, on skin, or on clothing.
- P264: Wash hands thoroughly after handling.
- P270: Do not eat, drink, or smoke when using this product.
- P391: Collect spillage.
- P280: Wear protective gloves, goggles, and protective mask.
- EUH 501: Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

Contains micro-organisms. May cause allergic reactions. The use of respiratory protection equipment, disposable respirator, or filter mask, at least N99 (EN 149 FFP3) or equivalent, is recommended when handling the preparation or applying the product.

First Aid Measures

Recommendations for Doctor. First aid measures general:

Never give anything by mouth to an unconscious person. If you feel unwell or experience respiratory symptoms, contact a doctor or call an anti-poison centre. If medical advice is needed, have product container or label at hand.

In case of inhalation: Bring the subject to fresh air. If breathing stops, give artificial respiration, Call a doctor.

In case of contact with skin: Remove contaminated clothes. Wash with plenty of water, If skin irritation occurs, contact a doctor for medical advice.

In case of contact with eyes: Rinse cautiously with water for several minutes keeping the eyes wide open. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, contact a doctor for medical advice.

If swallowed: Rinse mouth, Do NOT induce vomiting, Call a doctor or Poison centre.

If swallowed: Rinse mouth. Do NOT induce vomiting. Call a doctor or Poison centre immediately.

DO NOT LEAVE THE PATIENT ALONE.

Therapeutic advises: Symptomatic treatment

Call a doctor if you feel unwell. Have the product container or label at hand.

Environmental risk mitigation: DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water / avoid contamination via drains from farm yards and roads).

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



https://www.corteva.ie/content/dam/dpagco/corteva/eu/ie/en/

files/sds/BLUE-N-IRL-SDS.pdf

Alternatively contact your supplier

Blank area for in-house print of 2D data matrix code 5700483LBLFRT2211

BlueN[™]



NUTRIENT EFFICIENCY BIOSTIMULANT

BIOSTIMULANT

TO INCREASE NUTRIENT EFFICIENCY, UPTAKE AND AVAILABILITY.

Mode of Action: Increase nutrient availability, uptake and utilisation by fixation of atmospheric nitrogen (N_2) and making it directly available to the plant, and by increasing photosynthesis efficiency of the plant. BlueN $^{\text{TM}}$ contains an exclusive strain of Methylobacterium symbioticum SB23, a nitrogen fixing bacteria characterised by its ability to fix atmospheric nitrogen and transform it into a nitrogen source that can be assimilated by the crop. After application, the bacteria present in BlueN $^{\text{TM}}$ rapidly colonises the plant's phyllosphere and multibiles throughout the plant growth cycle.

BlueNTM provides a new natural nitrogen source to the crop through atmospheric nitrogen fixation and symbiotic interaction with a crop.

Authorised uses:

BlueN™ is intended for foliar application.

Crops*	Application Rate (g/ha)	Time of application**	Water volume I/ha
Arable crops – ap	ply when at lea	st 50% of the ground is covered by crops	
Winter cereals	333	Apply between mid of tillering (BBCH 25) and beginning of flowering (BBCH 61)	100-250
Spring cereals		Apply between mid of tillering (BBCH 25) and 2 nd node stage (BBCH 32)	100-250
Winter and spring oilseed rape		Apply at 6-8-leaf stage in autumn BBCH 16-18 and in spring from the rosette stage (BBCH 30) up to the end-of-flowering stage (BBCH 69)	100-250
Maize		Apply between 4-8 true leaves (BBCH 14-18)	100-300
Grain legumes		Apply between 4 and 8 leaf stage (BBCH 14 – 18)	100-250
Grassland		Apply between beginning of tillering stage (BBCH 21) and flag leaf stage (BBCH 39)	100-250
Potatoes		Apply between closing rows stage (BBCH 25) and beginning of flowering (BBCH 61)	100-400
Sugar beet		Apply between BBCH 14 – BBCH 39	100-400
Horticultural crops -	apply when at	least 50% of the ground is covered by crops	
Chard, garlic, artichoke, celery, broccoli, onion, green onions, cabbage, cauliflower, escarole, white asparagus, green asparagus, spinach, pea, broad bean, sweet beet, fennel, lettuce, leek, radish, and carrot, Lucerne	333	Apply between 4-leaf stage (BBCH 14) and beginning of flowering (BBCH 61)	100-400
Strawberry, blueberries, raspberries, blackberries, currants, cranberries		Apply between 5-leaf stage (BBCH 15) and beginning of flowering (BBCH 61)	100-400

^{*} For other crops and/or more information, consult with Corteva hotline.

Take care to avoid drift through the use of drift reduction technology and applying during periods of low wind speeds.

Recommendations for use

- Conditions of use

Slurry pH: Do not incorporate the product into a slurry with a pH less than 5 or more than 8. Observe a chlorine content of the slurry water of less than 2 ppm.

Apply in the morning to facilitate the penetration of the bacteria into the plant.

It is recommended to apply during the period of maximum opening of the stomata at temperatures between 10°C and 30°C.

Crop Condition: Apply during active crop growth.

- Precautions of use

Do not apply to crops under stress. Stress can be caused by many factors including drought, frost, ground compaction, water-logging, trace element deficiency, disease and pest attack etc.

Rainfastness: if there is a risk of rain, observe a minimum delay of more than 1 hour between the application of the product and the rain event. If these conditions are not met, re-apply the product.

- Compatibility

Do not apply when crops are under stress.

Application combined with chlorinated products or water (> 2 ppm CI-) is not recommended, nor surfactants, fertilisers or plant protection products that contain them.

BlueNTM has been tested in mixture with various plant protection products, fertilisers and adjuvants. For more information on compatibility, please contact Corteva Agriscience technical hotline.

Any mixtures should be applied within 6 hours after preparation.

Storage and disposal:

Keep the product only in its original packaging, away from food and drink, including those for animals. Keep out of the reach of children and unauthorised personnel.

Store the product at room temperature, in a well-ventilated area and in the original container. Do not expose to temperatures below 4°C or above 35°C. Keep the product tightly closed. After opening a pack, the product should be used quickly. Rinse the container thoroughly three times with clean water, making sure to pour the rinsing water into the sprayer tank. Empty packaging should be disposed of according to local requirements. Reuse of packaging is prohibited.

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in additional to the following personal protective equipment:WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling concentrate. WHEN USING DO NOT EAT, DRINK OR SMOKE.

In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show label where possible).

Sprayer cleaning

The entire sprayer (tank, boom, sump, circuit, nozzles, etc.) must be rinsed with clean water in accordance with the regulations in force.

- In the event of a spill

Wear suitable protective clothing, gloves, and eye or face protection. Secure the area. Collect anything that may have been in contact with the product.

Clean the site and the equipment used, taking care to contain the effluents generated by the cleaning operation. Dispose of them according to the regulations in force.

WARNING

Any total or partial reproduction of this label is prohibited.

The manufacturer offers this product under the following agreed conditions and sales guarantee:
The conditions of use of the product have been determined following rigorous tests and trials.
However, it is impossible to take into account all the variables and eliminate all the risks associated

with its use. Injury or damage may result from conditions beyond the control of the manufacturer. The manufacturer only guarantees that this product conforms to its composition appearing on the packaging and that it is considered to be reasonably suitable for the purposes referred to under normal conditions of use. Under no circumstances should the manufacturer be held responsible for indirect or specific damage resulting from the use or handling of the product. Any deviation from this warranty can only be valid if it is issued in writing by an authorised legal representative of the manufacturer.

Product registered in Spain under registration number F0004505/2030

Blank area for in-house print of 2D data matrix code

^{**} The BBCH growth stages are approximate and may vary depending on the weather conditions and agricultural practices.

Apply in the spring during periods of active growth.