

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection (UK only).

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking, and after work.

Environmental protection:

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from yards and roads.

Storage and disposal:

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

PROFESSIONAL USE ONLY

TRIPLE RINSE CONTAINER, PUNCTURE AND INVERT TO DRY AT TIME OF USE

**READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.
SHAKE WELL BEFORE USE.
PROTECT FROM FROST.**

Pack size: 5 Litres



CABADEX®

Product Registration Number: MAPP 13948/PCS No. 03749

A suspension emulsion formulation containing 100 g/litre fluroxypyr and 2.5 g/litre florasulam.

A post-emergence herbicide for use on MANAGED AMENITY TURF, including domestic lawns, and AMENITY GRASSLAND for the control of DAISY, DANDELION, CLOVER, BUTTERCUP, RIBWORT PLANTAIN and other broad-leaved weeds.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK only).



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008

[CLP]: Cabadex®; Hydrocarbons, O9, aromatics

WARNING

CAUSES SKIN IRRITATION

MAY CAUSE AN ALLERGIC SKIN REACTION

CAUSES SERIOUS EYE IRRITATION

MAY CAUSE RESPIRATORY IRRITATION

MAY CAUSE DROWSINESS OR DIZZINESS

VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS

Wear protective gloves/clothing/eye/face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing. Do NOT induce vomiting.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean triple rinsed containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP 13948/PCS No. 03749

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL HERBICIDE

| Situation | Maximum Individual Dose | Maximum Number of Applications |
|--|--------------------------------|--------------------------------|
| Managed amenity turf, lawns, amenity grassland | 2.0 litres product per hectare | One per year |

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for using Plant Protection Products.


Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE.

Telephone: +44 (0) 1462 457272 Fax: +44 (0) 1462 426605

For Technical Enquiries Call: +44 (0) 800 6898899

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251



®Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow

This label is compliant with the CPA Voluntary Initiative Guidance (UK only).



Headland Amenity Limited

1010 Cambourne Business Park, Cambourne, Cambridgeshire, CB23 6DP Telephone: +44 (0) 1223 597834
24 Hour Emergency Telephone Number: +44 (0) 1553 761 251

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

CABADEX® herbicide has activity against a range of broad-leaved weeds. The ideal timing for application is when the weeds are small and actively growing.

NOTES

Broad-leaved weeds not present at application will not be controlled.

Clippings from grass treated with CABADEX can be safely used for mulch after the third cut.

An interval of four weeks must elapse between application of CABADEX and re-seeding turf.

Do not apply if turfgrass is wet.

Do not apply to turf, lawns or grass areas which are under stress.

Do not apply if night temperatures are low, if ground frost is imminent, or in periods of prolonged cold or dry weather.

Ensure weeds are actively growing as after periods of prolonged drought, weeds can take a long time to start actively growing again after soil moisture returns.

Take extreme care to avoid drift onto crops and non-target plants e.g. trees, shrubs, bedding, outside the target area.

RESISTANCE

CABADEX contains active ingredients with differing modes of action and the risk of resistance building is therefore reduced. However, as florasulam is an ALS-inhibitor there is a risk of resistance building to this active ingredient and so precautions should be taken to minimise the risk. Therefore, avoid using single action mode of action herbicides, such as ALS-inhibitors in the same field over a number of years. Users are advised to apply products containing herbicides with different modes of action or use sequences or tank mixtures where two or more components are active against the target weeds.

AREA OF USE

CABADEX can be applied to newly sown or established managed amenity turf, including domestic lawns, and amenity grassland.

Ensure newly sown turf has become established before treating. Turf sown in spring or summer may be ready for spraying at or after two leaf growth stage, usually two months after sowing, but turf sown in late summer or autumn should not be sprayed until growth is resumed in the following spring, perhaps eight months after sowing.

CABADEX has been tested for selectivity on the following range of turf grass species:

| | |
|---------------------|-----------------------------|
| Annual meadow-grass | Perennial ryegrass |
| Browntop bent | Smooth-stalked meadow-grass |
| Chewings fescue | Rough stalked bluegrass |
| Creeping bentgrass | |

In view of the large number of turf grass cultivars grown consult manufacturer for current approved list or test CABADEX for turf safety on a small area of turf before overall application.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

APPLICATION TIMING

Apply when weeds are in active growth normally from March to October when the soil is moist. Do not apply in periods of drought unless irrigation is applied. Avoid mowing 3 days before and after spraying to ensure sufficient weed leaf surface is present and to allow uptake and movement of CABADEX within the weed.

RATE OF APPLICATION AND WEEDS CONTROLLED

One application of CABADEX will control susceptible emerged weeds at the following rates:

| Weed | Rate L product/ha | Rate ml product/100m ² |
|----------------------------------|-------------------|-----------------------------------|
| Bird's-foot trefoil ¹ | 2.0 | 20 |
| Common daisy | | |
| Common dandelion | | |
| Common mouse-ear | | |
| Creeping buttercup | | |
| Ribwort plantain ¹ | | |
| Slender speedwell | | |
| White clover | | |
| Yarrow ¹ | | |

¹Moderate control only

WATER VOLUME

For overall application apply CABADEX in 200 litres of water per hectare. For knapsack application apply CABADEX in 2 litres of water per 100 m².

APPLICATION EQUIPMENT

CABADEX may be applied through tractor-mounted hydraulic sprayers or knapsack sprayers providing they are in good working order and have been calibrated according to the manufacturers' recommendations.

Do not apply through CDA applicators.

MIXING

Half fill the spray tank with water and add the required amount of CABADEX. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing.

SPRAY QUALITY

Apply CABADEX as a MEDIUM spray as defined by the BCPC system.

TANK CLEANING

To avoid subsequent injury to crops other than managed amenity turf, domestic lawns and amenity grassland, all spraying equipment must be thoroughly cleaned both inside and out, using All Clear Extra spray cleaner as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water and add All Clear Extra at the recommended rate. Agitate and then briefly flush the booms and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Flush the booms and hoses and drain tank completely.
4. Nozzles and filters should be removed and cleaned separately with All Clear Extra solution containing 50 ml of All Clear Extra per 10 litres of water.
5. Rinse the tank with clean water and flush through the booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
6. For disposal of washings, follow The Code of Practice for Using Plant Protection Products. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

Dow AgroSciences Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:

UK



<http://uk.dowagro.com/wp-content/uploads/CABADEx-SDS.pdf>

Alternatively contact your supplier

REPUBLIC OF IRELAND

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<http://uk.dowagro.com/wp-content/uploads/CABADEx-IRL-SDS.pdf>

Alternatively contact your supplier

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection (UK only).

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking, and after work.

Environmental protection:

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from yards and roads.

Storage and disposal:

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

PROFESSIONAL USE ONLY

TRIPLE RINSE CONTAINER, PUNCTURE AND INVERT TO DRY AT TIME OF USE

**READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.
SHAKE WELL BEFORE USE.
PROTECT FROM FROST.**

Pack size: 5 Litres




CABADEX®

Product Registration Number: MAPP 13948/PCS No. 03749

A suspension emulsion formulation containing 100 g/litre fluroxypyr and 2.5 g/litre florasulam.

A post-emergence herbicide for use on MANAGED AMENITY TURF, including domestic lawns, and AMENITY GRASSLAND for the control of DAISY, DANDELION, CLOVER, BUTTERCUP, RIBWORT PLANTAIN and other broad-leaved weeds.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK only).



Product Identifier according to Art. 18 of Reg. (EC) No 1272/2008

[CLP]: Cabadex®; Hydrocarbons, C9, aromatics

WARNING

CAUSES SKIN IRRITATION

MAY CAUSE AN ALLERGIC SKIN REACTION

CAUSES SERIOUS EYE IRRITATION

MAY CAUSE RESPIRATORY IRRITATION

MAY CAUSE DROWSINESS OR DIZZINESS

VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS

Wear protective gloves/clothing/eye/face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing. Do NOT induce vomiting.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean triple rinsed containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP 13948/PCS No. 03749

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL HERBICIDE

| Situation | Maximum Individual Dose | Maximum Number of Applications |
|--|--------------------------------|--------------------------------|
| Managed amenity turf, lawns, amenity grassland | 2.0 litres product per hectare | One per year |

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for using Plant Protection Products.


Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE.

Telephone: +44 (0) 1462 457272 Fax: +44 (0) 1462 426605

For Technical Enquiries Call: +44 (0) 800 6898899

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251



*Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow

This label is compliant with the CPA Voluntary Initiative Guidance (UK only).



Headland Amenity Limited

1010 Cambourne Business Park, Cambourne, Cambridgeshire, CB23 6DP Telephone: +44 (0) 1223 597834

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251