

Curzate[®] M WG

FUNGICIDE


GROUP

M3;27

FUNGICIDE

PCS No.: 04723

A water dispersible granule containing 4.5% w/w cymoxanil and 68% w/w mancozeb for the control of potato late blight in po



Avoid breathing vapours/spr Wear protective al If medical advice el at hand

> CTIONS F Y PRECAI ROTECTION

Do not concaminate water with th surface water/Avoid contamination and roads). t aquatic organisms respe RICTIONS

• Do not kee,

Keep away from

ATE M WG cor moxanil and the protectant activity of mancozeb

same mode of action are used repeatedly over several years in the same field, s sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with differing modes

Contaminated work clothing should not be allowed out of the workplace IF ON SKIN: Wash with plenty of soap and water. IF exposed or concerned: Get medical advice/attention. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before Keep out of reach of children Store locked up Collect spillar

Dispos ept for t ks to human health and ti instr for use.

74723

CURZATE M grown for pro

D WEATHER Timing of spray app ant the normal prevalence of the disease in the area and bray should be applied as soon as there is a risk of blight ons in the par sued. In the absence of a warning, spraying should begin just

should be repeated at 10-14 day intervals depending on the prevalence of the disease and trable conditions for infection. A 7 day spray interval may be used under exceptional circumstances e.g. blight risk is already high and rain is forecast at the time of the next 10 day spray is due. The longer interval is recommended under dry conditions. Irrigation of crops increases the risk of blight and crops should always be sprayed with CURZATE M WG as soon as possible after irrigation subject to the minimum spray interval. High-risk situations include conditions of rapid growth, irrigated crops, areas of intensive potato growing, particularly blight-sensitive varieties, when further warnings are issued, or when active blight is in the area.

Apply CURZATE M WG at 2.3 kg/ha

DOSE

VOLUME AND APPLICATION Apply CURZATE M WG in not less than 200 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC). FOR USE ON JULTURAL

l l	N d	c e	Max no of applications	Max dose ('.at. oplic	of
Po	ed 2.	 .3kg		13.8kg/h	lays b	٠ ٦

ctor mounted / traile

IS PRUDUCT IN A ANNE IS INC THE LABEL MAY BE I OFFE OLLOW __ CUDE OF PRACTICE FOR USING ON PRODUCTS. ANT PI

WARNING

KEEP AWAY FROM FIRE OR SPARKS. Do not allow CURZATE M WG to become wet during storage or keep opened packs from one season to another as this may reduce the effectiveness of the fungicide and may create vapours that may be flammable.

NOTICE TO BUYER

All goods supplied by us are of high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

Du Pont (UK) Limited, Crop Protection Products Kings Court, London Road, Stevenage, Herts SG1 2NG, England Email: ukhotline@corteva.com Tel: +44 (0) 800 689 8899 Emergency Tel: (01) 901 4670 Tel (01) 809 2166

Manufactured in EU

FOR PROFESSIONAL USE ONLY

 ${}^{\circledR} \text{Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.}$

Contents:





22/10/2019 13:36



5H4/Y11/S/19 B/1103.2028-130204

Batch No: See on the pack



CURZATE M68 WG IF 10KG BAG K-44763 indd 1 P.186 P.285 BLACK P. YELLOW P.514 P.513 SPECINI